



Research Article

Analgesic, CNS depressant and antibacterial activities of ethanol extract of *Spilanthes paniculata* leaves

Manuscript received: 10 April 2024

Revision received: 16 May 2024

Accepted: 20 May 2024

Published: 21 May 2024

Md. Khalequeuzzaman*, Sadia Islam, Mst. Mahfuza Khatoon,
Anik Kumar Dey

Edited by:

Dr. Sreemoy Kanti Das

Faculty of Pharmacy

Lincoln University College, Malaysia

Citation:

Khalequeuzzaman M, Islam S, Khatoon MM, Dey AK. Analgesic, CNS depressant and antibacterial activities of ethanol extract of *Spilanthes paniculata* leaves. Healthmed J Pharm Sci 2024; 2(2):01-06.

**Copyright:** © 2024 by the authors.

Licensee: HMS Media

International, Dhaka, Bangladesh

This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution 4.0 (CC BY 4.0) International license.

Abstract

Spilanthes paniculata Wall. is an important medicinal plant which is used to treat toothache, throat and gum infections. The present study was aimed to investigate the analgesic, CNS depressant and antibacterial activities of the ethanol extract of *Spilanthes paniculata* leaves (SPE). Analgesic activity was determined by formalin and acetic acid induced writhing methods. The CNS depressant activity was determined by open field and hole cross test. Antimicrobial activity was evaluated by disk diffusion method. In acetic acid induced writhing method, the SPE showed significant writhing inhibition 66.92% ($p < 0.01$) at a dose of 500 mg/kg body weight as compared to control. In formalin test, the SPE (500mg/kg) showed a significant licking inhibition 46.77% ($p < 0.01$) In CNS depressant activity, test animals showed significant decrease ($p < 0.001$) in number of movement with the treatment of SPE (500mg/kg) while diazepam (1mg/kg) was used as a standard drug. SPE at the dose of 800 μ g/disc showed moderate antibacterial activity against most of the gram positive and gram-negative bacteria. Thus, our study suggests that ethanol extract of *Spilanthes paniculata* may be a source of natural product having its analgesic and CNS depressant activities along with moderate antibacterial activity.

Keywords:

Spilanthes paniculata, Ethanol extract, Analgesic, CNS depressant, Phytomedicine, Natural product, Antibacterial activity

Department of Pharmacy, Gono Bishwabidyalay (University), Nolam, Mirzanagar, Savar, Dhaka 1342, Bangladesh

***Correspondence:** Md. Khalequeuzzaman, Associate Professor, Department of Pharmacy, Gono Bishwabidyalay (University), Nolam, Mirzanagar, Savar, Dhaka 1342, Bangladesh; Email: sobujph@gmail.com; Phone: +8801924865471

Introduction

The empirical use of plants as medicine can be traced back to over five millennia to ancient documents to early civilizations, such as in China, Egypt, India and the near east, but is certainly as old as mankind (Balick and Cox, 2020). Many of the plants used in antiquity as medicines still play an important role for health today (Miraldi and Bains, 2019). The use of the medicinal plant is increasing in many countries where one third of drugs contain natural products. *Spilanthes paniculata* belongs to Asteraceae family is a small tender annual that grows to about 12-15 inches and will spread to 24-30 inches which is native to the Americas and has been introduced to Asia, Africa, the Pacific islands, and Australia. The most common and widespread medicinal use for *Spilanthes paniculata* is to treat toothache, throat and gum infections. A mouth rinse of spilanthes extract can be used daily to promote gum health, and chewing as little as a single bud of the plant can numb the mouth and reduce the pain of toothache for up to 20 minutes depending on the sensitivity of the person (Pathak et al 2013). The leaves of *Spilanthes paniculata* possesses antioxidant activity (Haque et al, 2015) and also have antidiabetic and thrombolytic effects (Akter et al, 2014).

The role of free radicals can be found in the inflammatory process which is a complex process resulting many human diseases (Biswas et al, 2017). Pain is formally defined as an unpleasant sensory and emotional incident coupled with real or likely tissue injury. Pain acts as a word of warning sign against disorder of the body and has a practical function.

Analgesic mitigates pain as a symptom without affecting its reason (Akter et al, 2009). However, data from multiple placebo-controlled trials and meta-analyses studies alarmingly signify the adverse-effects of NSAIDs in gastrointestinal, cardiovascular, hepatic, renal, cerebral and pulmonary complications (Bindu et al, 2020). As a result, more and more people are turning to herbal medicines as the alternative treatment of pain.

Anxiety disorders are often comorbid with one another and with other mental disorders, especially depression, as well as with somatic disorders. Such comorbidity generally signifies more severe symptoms, greater clinical burden, and greater treatment difficulty (Penninx et al, 2021). Notably, benzodiazepines and drugs based on neurotransmitter systems are currently being used to treat anxiety. However, ineffective treatment, undesirable effects and dependence still exist in many cases. The CNS depressant effect of herbs has been paid more and more attention gradually because of increasing incidence of anxiety and predominance of traditional herbs in therapy.

Antimicrobial agents are essentially important in reducing the global burden of infectious diseases (Bhatia et al 2010). However, emergence and dissemination of multidrug resistant (MDR) strain in pathogenic bacteria have become a significant public health threat as there are fewer, or even sometimes no, effective antimicrobial agents available for the infection caused by pathogenic bacteria (Giamarellou et al 2010). A vast number of medicinal plants have been recognized as valuable resources of natural antimicrobial

compounds as an alternative that can potentially be effective in the treatment of these problematic bacterial infections.

In order to explore the potential biological activity of the ethanol extract of *Spilanthes paniculata* leaves we studied analgesic, CNS depressant and antibacterial activity.

Materials and Methods

Chemicals

All the chemicals and reagents were used for this study was analytical grade from Merck Germany and Sigma Aldrich (USA). The standard drug Diazepam was collected from the Square Pharmaceuticals Ltd, Bangladesh.

Plant Material

The leaves of *Spilanthes paniculata* were collected from Dhamrai, Dhaka, Bangladesh and was identified at the department of Botany, Jahangirnagar University, Savar, Dhaka, Bangladesh. Immediately after collection, the leaves were thoroughly washed with water. Then the leaves were dried under shade for 2 days and were ground to coarse powder with a mechanical grinder. The powder was extracted individually with ethanol and ether in a Soxhlet apparatus. The mixture was filtered and the filtrate was concentrated by Rota evaporator to yield semisolid mass. The extracts were preserved in refrigerator until further use.

Experimental animals

Swiss albino mice of both sexes, aged 5-6 weeks, weight about 20-30 g were purchased from the department of Pharmacy of Jahangirnagar University, Savar, Dhaka, Bangladesh. Before initiating the experiment, the mice were kept in standard environmental conditions having temperature at $22.0 \pm 2.0^\circ\text{C}$, relative humidity at 55-65% and 12 h light/12 h dark cycle and had free access to feed and water ad libitum. The animals were acclimatized to laboratory condition for one week prior to experiments.

Ethical approval of experiment protocol

The ethical review board of Gono University in Bangladesh approved the protocol for animal experiments.

Determination of analgesic activity by acetic acid-induced writhing test

The acetic acid-induced writhing method is an analgesic behavioral observation assessment method that demonstrates noxious stimulation in mice. The experiment was carried out following the method described by Sharma et al 2010. Fifty mg of crude ethanol extracts were triturated by adding a small amount of suspending agent (Tween 80). Normal saline (0.9% NaCl) was slowly added to make the final volume up to 2.5ml. To prepare the standard, indomethacin 10 mg was dissolved into 0.9% normal saline and increased the volume to 10 ml. The control solution, tween 80 (1%) was mixed properly in the normal saline to make the volume up to 5 ml. The test samples, control (1% tween 80 in water) and standard (indomethacin) were administered orally with the help of a feeding needle at the beginning of the experiment. After 30 minutes, 0.7% acetic acid was injected intra-peritoneally to each of the animals of all the groups to create a pain sensation. Then the animals were placed on an observation table. Approximately 5 minutes after the injection of acetic acid, a wave of contraction and elongation of abdominal musculature referred to as writhing is started and the number of writhing

for the next 10 minutes was counted for each rat. Full writhing was not always accomplished by the animal, because sometimes the animals started to gibe writhing but they did not complete it. This incomplete writhing is considered as half-writhing. Accordingly, two half-writhing was considered as one full writhing. The number of writhes in each treated group was compared to that of a control group while indomethacin (10 mg/kg) was used as a reference substance (positive control). The percent inhibition (% analgesic activity) was calculated by the following equation:

$$\% \text{ inhibition} = [(A-B)/A] \times 100$$

where,

A= Average number of writhing of control group

B= Average number of writhing of test group

Determination of analgesic activity by formalin-induced hind paw-licking test

The analgesic activity of the drugs was determined using the formalin test as described by Sharma et al, 2010. Fifty mg of crude ethanol extract of *Spilanthes paniculata* was triturated by the addition of small amount of suspending agent. Normal saline (0.9% NaCl) was slowly added to make the final volume up to 2.5 ml. To prepare the standard, indomethacin 10 mg was dissolved into 0.9% normal saline and made the volume up to 10 ml. For preparing control sample, distilled water was mixed properly in the normal saline to make the volume up to 5 ml. The test samples, control (distilled water 10 ml/kg) and standard (indomethacin 10 mg/kg) are administered orally with the help of a feeding needle at the beginning of the experiment. After 30 minutes, 0.05ml of 2.5% formalin (40% formaldehyde) in distilled water is injected into the dorsal surface of the right hind paw of the mice. Then the mice are individually placed in transparent cage observation chamber. The time spent licking the injected paw is recorded and the data were expressed as total licking time in the early phase (0-5 min) and the late phase (15-30 min) after formalin injection. The percent inhibition (%licking activity) is calculated by the following equation:

$$\% \text{ inhibition} = [(A-B)/A] \times 100$$

Where, A = Average number of licking of the control per group

B = Average number of licking of the test per group

CNS depressant activity by hole cross method

The most consistent behavioral change is a hyperemotional response to a novel environment. The experiment was carried out as described by Munira et al, 2019. The experiment was carried out to characterize the emotional behavior of mice using the hole-board test. The mice are divided into control, standard control and test group. The test groups receive ethanol extract of leaves of *Spilanthes paniculata* at the dose of 250 and 500 mg/kg body weight orally whereas control group receive vehicle (1% Tween 80 in water) at 10ml/kg body weight orally and standard group receive diazepam at the dose of 1 mg/kg body weight orally with the help of a feeding needle at the beginning of the experiment. A steel partition is fixed in the middle of a cage having a size of 30×20×14 cm. A hole of 3cm diameter is made at a height of 7.5cm in the center of the cage. The number of passages of a rat through the hole from one chamber to other is counted for a period of 3 min on 0, 30, 60, 90 and 120min after oral administration of test drugs.

CNS depressant activity by open field method

This experiment was carried out as described by Sultana et al, 2018. The open field test is clearly the most frequently used of all behavioral tests in pharmacology and neuroscience. The mice are divided into control, standard and test group. The test groups receive ethanol extract of leaves of *Spilanthes paniculata* at the dose of 250 and 500 mg/kg body weight orally whereas control group receive vehicle (1% Tween 80 in water) at 10ml/kg body weight orally and standard group received diazepam at the dose of 1mg/kg body weight orally with the help of a feeding needle at the beginning of the experiment. The floor of an open field of half square meter is divided into a series of squares each alternatively colored black and white. The apparatus has 40 cm height a wall. The number of squares visited by the animals was counted for 3 min, on 0, 30, 60, 90 and 120 min after oral administration of standard and sample.

Anti-bacterial Activity

Antibacterial activity of the ethanol extracts of leaves of *Spilanthes paniculata* was determined by disc diffusion technique (Ngamsurach and Praipipat, 2022). To determine the antibacterial activity of ethanol extract two gram-positive (*Staphylococcus aureus*, *Bacillus cereus*) and two gram-negative (*Escherichia coli*, *Pseudomonas aeruginosa*,) bacteria were used. For comparison kanamycin-K (30 µg per disc) was used as standard. Nutrient agar media was reconstituted with distilled water in a conical flask according to specification (2.3%). It was then heated in water bath to dissolve the agar until a clear solution of agar was obtained. The media prepared was then transferred in 20ml and 5ml to prepare plates and slants respectively in a number of clean test tubes. The slants were used for making sub-culture of microorganism which in turn were used for sensitivity tests. The tubes were then plugged with cotton and sterilized in an autoclave at a temperature of 121°C and a pressure of 151 lb/sq. inch for 20 minutes. The test organisms were transferred from the pure culture to the agar slants with the help of an inoculating loop in an aseptic condition. The inoculated slants were then incubated at 37°C for 18-24 hours to assure the growth of the test organisms. This culture was then used within one week. With the help of an inoculating loop, the test organism was transferred from the subculture to the test tube containing 20 ml autoclaved media in an aseptic area. Then The test tube was shaken well by rotation to get a uniform suspension of organism. The bacterial suspensions were immediately transferred to the sterile petri-dishes in such a way as to obtain a uniform depth of media (approximately 4mm thick). The petri-dishes were rotated several times, first clockwise and then anti-clockwise to assure homogenous distribution of the test organisms. The plates were cooled to room temperature and it was stored in a refrigerator (4°C). After that the test sample and standard discs were prepared carefully at concentration of 800µg/disc and 30µgm/discs respectively. The Prepared discs were placed gently on the freshly seeded solidified agar plates with a sterile forceps. Standard discs and controlled discs were also placed on the test plates. The spatial arrangements of the discs were such that the discs were not closer than 15mm to the edge of the plates to prevent overlapping the zone of inhibition. The plates were then inverted and kept in a refrigerator for about 1244 hours at 4°C to obtain maximum diffusion. Finally, the plates were incubated at 37°C for 12-18 hours. After incubation, the

antibacterial activities of the test samples were determined by measuring the diameter of inhibitory zones in terms of millimeters (mm).

Results

Analgesic activity of *Spilanthes paniculata* leaves by Acetic acid-induced writhing test

In acetic acid-induced writhing test, the ethanol extract of *Spilanthes paniculata* leaves significantly and dose dependently suppressed the frequency of acetic acid-induced writhing in mice after oral administration. At 500 mg/kg body weight, SPE showed 66.92% of writhing inhibition whereas at 10 mg/kg body weight, the standard drug indomethacin showed 60% of writhing inhibition.

Table 1: Analgesic activity of *Spilanthes paniculata* leaves by Acetic acid-induced writhing test

Treatment Group	Dose (body weight)	Mean no. of writhing	% of inhibition
1% Tween 80 in water (control)	1 ml /10 gm	32.50±8.74	-
Indomethacin (standard)	10 mg/kg	13.00±3.65**	60
SPE	250 mg/kg	16.50±5.19*	49.23
	500 mg/kg	10.75±2.99**	66.92

All the values are stated as Mean ± SD. (Where, n=4); significance level ***p<0.001, **p<0.01, *p<0.05 compared to control.




Fig 1. Analgesic activity of *Spilanthes paniculata* leaves by Acetic acid-induced writhing test

Analgesic activity of *Spilanthes paniculata* leaves by Formalin-induced pain test

Table 2. Analgesic activity of *Spilanthes paniculata* leaves by formalin-induced pain test

Treatment Group	Dose	Mean number of lickings	% of inhibition
Distilled water (control)	10 ml/kg	31.00±7.96	-
Indomethacin (standard)	10 mg/kg	19.50±1.29*	37.10
SPE	200 mg/kg	21.50±2.08*	30.64
	500 mg/kg	16.50±3.42**	46.77

All the values are Mean ± SD. (Where n=4); significance level ***p<0.001, **p<0.01, *p<0.05 compared to control.

The ethanol extract of leaves of *Spilanthes paniculata* significantly suppressed the licking activity in the formalin-induced pain in mice in a dose-dependent manner. The standard drug (Indomethacin) used in the experiment showed 37.10% licking inhibition with a dose of 10 mg/kg of body weight whereas SPE showed 46.77% licking inhibition with a dose of 500 mg/kg of body weight.




Fig 2. Analgesic activity of *Spilanthes paniculata* leaves by formalin-induced pain test

CNS depressant activity of *Spilanthes paniculata* leaves by hole cross test

The animal treated with different doses of ethanol extract of *Spilanthes paniculata* leaves showed a dose-dependent reduction of locomotor activity that was comparable with that of standard drug diazepam. The extract reduced spontaneous motor activity, and this effect may be attributed to CNS depression, as depression of locomotor activity is common to most neuroleptics.

Table 3. CNS depressant activity of *Spilanthes paniculata* leaves by hole cross test

Treatment Group	Dose (body weight)	No. of movements				
		0 min	30 min	60 min	90 min	120 min
1% tween 80 in water (control)	10ml/kg	12.75±0.96	13.50±2.38	13.75±2.21	9.25±4.57	12.00±2.58
Diazepam (standard)	1 mg/kg	9.25±0.96*	7.75±1.25*	6.75±1.26*	4.25±1.50	3.00±1.41**
SPE	250 mg/kg	11.25±0.96	10.75±1.89	7.75±0.96	6.75±1.26	5.75±0.50*
	500 mg/kg	9.50±0.96***	4.25±0.96***	2.75±0.96***	2.00±0.82**	1.50±0.57***

All the values are stated as Means ± SD (Where, n=4); significance at ***p<0.001, **p<0.01, *p<0.05 as compared to control.

CNS depressant activity of *Spilanthes paniculata* leaves by open field test

Open-field test was carried out to determine the depressive action of the test samples. In this test, the extract showed a noticeable dose dependent decrease in locomotion in the test animals. Test animals showed significant decrease in number of movements at a dose of 500 mg/kg after 120 min compared to standard.

Table 4. CNS depressant activity of *Spilanthes paniculata* leaves by Open field test

Treatment Group	Dose (body weight)	No. of movements				
		0 min	30 min	60 min	90 min	120 min
1% tween 80 in water (control)	10ml/kg	62.50 ±9.34	57.50 ±6.75	55.25 ±6.23	51.75 ±4.03	49.00 ±5.47
Diazepam (standard)	1 mg/kg	55.00 ±7.26	25.00 ±4.96 ***	12.25 ±2.98 ***	10.50 ±2.38 ***	5.00± 2.61* **
SPE	250 mg/kg	82.75 ±3.77	74.00 ±5.97 **	58.50 ±8.10	47.25 ±7.88	34.25 ±6.34 *
	500 mg/kg	58.25 ±7.50	28.50 ±2.88 ***	15.75 ±2.06 ***	9.75± 1.71* **	4.75± 0.96* **

All the values are stated as Mean ± SD. (Where, n=4); significance at ***p<0.001, **p<0.01, *p<0.05 as compared to control.

Anti-bacterial activity

Evaluation of the antimicrobial activity of the ethanol extract of *Spilanthes paniculata* was determined initially by the disc diffusion method against different microorganisms. These organisms were frequently encountered in infectious diseases. It was observed that the extract used in the study exhibited a varying degree of antibacterial activity against all microorganisms tested.

Table 5. Antibacterial activity of *Spilanthes paniculata* leaves by disk diffusion method.

Bacteria		Diameter of the zone of inhibition (mm)	
		SPE 800 µg/disc.	Kanamycin-K 30 µg/disc.
Gram Positive	<i>Staphylococcus aureus</i>	16	25
	<i>Bacillus cereus</i>	9	26
Gram Negative	<i>Pseudomonas aeruginosa</i>	6	20
	<i>Esherichia coli</i>	8	22

Discussion

Scientific and methodical investigation of herbal plants has become a potential source for the discovery of lead compounds of high therapeutic value in terms of analgesic activity. Ethno-pharmacological studies have become increasingly invaluable in the development of modalities for the management of pain and related disorders. Thus green pharmaceuticals have now received considerable attention and popularity in this area due to its availability, less side effects and economic feasibility compared to the orthodox medicine. In this study the ethanol extract of the *Spilanthes paniculata* leaves showed significant analgesic activity (p<0.01) at a dose of 500 mg/kg body weight compared to standard indomethacin. So, the leaves of this plant can be used as an alternative analgesic medicine.

Central nervous system (CNS) depressants are drugs that can be used to slow down brain activity. CNS depressants may be prescribed by a physician to treat anxiety, muscle tension, pain, insomnia, acute stress reactions, panic attacks, and seizure disorders. CNS depression often results from the use of depressant drugs such as alcohol, opioids, barbiturates, benzodiazepines, general anesthetics etc. Drug overdose is often caused by combining two or more depressant drugs, although overdose is certainly possible by consuming a large dose of one depressant drug. In this study the ethanol extract of the *Spilanthes paniculata* leaves showed significant CNS depressant activity (p<0.001) at a dose of 500 mg/kg body weight compared to standard diazepam. So the leaves of this plant can be used as an alternative CNS depressant medicine. In the light of the evidence of the rapid global spread of antibiotic resistant bacterial strains, the need to find new antimicrobial agents is of paramount importance. Plants are rich in a wide variety of secondary metabolites such as tannins, alkaloids, phenolic compounds, and flavonoids, which have been found in vitro to have antimicrobial properties (Duraipandiyar et al, 2006). In this study the ethanol extract of the *Spilanthes paniculata* leaves showed moderate antibacterial activity against *Staphylococcus aureus* among the four bacterial strains compared to standard Kanamycin.

Conclusion and Future Direction

This study revealed that the leaf extract of *Spilanthes paniculata* possesses good analgesic and CNS depressant activity. The plant extract also has moderate antibacterial activity. Further more specific studies may confirm the analgesic and CNS depressant potential of this plant which could open a new window on the use of this plant in traditional medicine.

List of Abbreviations

SPE: Ethanol extract of *Spilanthes paniculata*, SD: Standard Deviation, CNS: Central Nervous System.

Data Availability Statement

Data relevant to the study is already included to the article or attached in the supplements. Raw data will be provided on reasonable request upon contacting with the corresponding.

Authors Contributions

M. Khalequeuzzaman and M. Khatoun designed the study protocol and S. Islam collected the plant. M. Khalequeuzzaman, M. Khatoun, and S. Islam performed the experiments. Anik Kumar Dey contributed to data analysis.

Conflicts of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Funding Information

This research did not receive any funding or grant support.

Acknowledgments

Authors are grateful to the department of Pharmacy, Gono Bishwabidyalay in where this research work was conducted.

References

- Akter R, Hasan SMR, Siddiqua SA, Majumder MM, Hossain MM, Alam MA, et al. Evaluation of analgesic and antioxidant potential of the leaves of *Curcuma alismatifolia* Gagnep. S. J. Pharm. Sci; 2009 ; 1&2: 3-9. DOI: <http://doi.org/10.3329/sjps.v1i1.1779>.
- Akter S, Rahman MA, Azad MAK, Mohiuddin M, Mamun AA, Sarker J et al. Antidiabetic and thrombolytic effects of ethanol extract of *Spilanthes paniculata* leaves. Journal of Plant Sciences; 2014; 2(6-1): 13-18. DOI: 10.11648/j.jps.s.2014020601.13.
- Balick MJ, Cox PA. Plants, People, and Culture: the Science of Ethnobotany, Scientific American Library; 2020; New York, U.S.A. DOI: 10.4324/9781003049074.
- Bhatia R and Narain JP. The growing challenge of antimicrobial resistance in the South-East Asia Region - are we losing the battle. *Indian Journal of Medical Research*; 2010; 132(5), 482–486. DOI:10.4103/0971-5916.73313.
- Bindu S, Mazumder S, Bandyopadhyay U. Non-steroidal anti-inflammatory drugs (NSAIDs) and organ damage: A current perspective. *Biochem Pharmacol*; 2020;1;180. DOI: 10.1016/j.bcp.2020.114147.
- Biswas S, Das R and Banerjee ER. Role of free radicals in human inflammatory diseases. *AIMS Biophysics*; 2017; 4(4): 596-614. DOI: 10.3934/biophy.2017.4.596.
- Giamarellou H. Multidrug-resistant Gram-negative bacteria: how to treat and for how long. *International Journal of Antimicrobial Agents*; 2010; 36 (Suppl 2): S50–S54. DOI: <https://doi.org/10.1016/j.ijantimicag.2010.11.14>.
- Haque S, Lopa SD, Das BK. Evaluation of antioxidant potential of *Spilanthes paniculata*. *Int J Pharm Pharm Sci*; 2015; 7(8): 390-392. DOI: <https://journals.innovareacademics/6703/6893>.
- Miraldi M and Bains G. Medicinal plants and health in human history: from empirical use to modern phytotherapy. *Journal of the Siena Academy of Sciences*; 2019, 10(1). DOI: 10.4081/jsas.2018.8529.
- Munira MS, Ahmed SN, Islam MS, Nesa ML and Kabir MH. Pharmacological Screening for CNS Depression, Analgesic and Anti-inflammatory Potentials of *Sonneratia caseolaris* (Linn.) Barks in Different Solvent Fraction. *Journal of Pharmaceutical Research International*; 2019; 29(5); 1-11. DOI: 10.9734/jpri/2019/v29i530251.
- Ngamsurach P and Praipipat P. Antibacterial activities against *Staphylococcus aureus* and *Escherichia coli* of extracted Piper betle leaf materials by disc diffusion assay and batch experiments. *ROYAL SOCIETY OF CHEMISTRY*; 2022, 12, 435-454. DOI: 10.1039/D2RA04611C.
- Pathak K. Herbal medicine- A Rational Approach in Health Care System. *International Journal of Herbal Medicine*. 2013; 1(3): 86-89.
- Penninx BW, Pine DS, Holmes EA and Reif A. Anxiety disorders. *The Lancet*; 2021; 397(10277), 914-927. DOI: 10.1016/S0140-6736(21)00359-7
- Sharma A, Bhatial S, Kharyaz MD, Gajbhiye V, Ganesh N, Namdeo AG et al. Anti-inflammatory and analgesic activity of different fractions of *Boswellia serrata*. *Int J Phytomed*; 2010; 2: 94-99. DOI: 10.5138/ijpm.2010.0975.0185.02015.
- Sultana T, Mannan MA, and Ahmed T. Evaluation of central nervous system (CNS) depressant activity of methanolic extract of *Commelina diffusa* Burm. in mice. *International Journal of Phytomedicine and Phytotherapy*; 2018; 4(5). DOI: 10.1186/s40816-018-0063-1.